JOURNAL OF CHROMATOGRAPHY A

VOL. 763, NOS. 1,2 28 FEBRUARY 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

20TH INTERNATIONAL SYMPOSIUM ON HIGH PERFORMANCE LIQUID PHASE SEPARATIONS AND RELATED TECHNIQUES, SAN FRANCISCO, CA, USA, 16–21 JUNE 1996, PART II

Applications of proteins and their constitutents

Tryptophan analysis in peptides and proteins, mainly by liquid chromatography (Review)
by I. Molnár-Perl (Budapest, Hungary) 1

Using quaternary high-performance liquid chromatography eluent systems for separating 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate-derivatized amino acid mixtures
by C. van Wandelen and S.A. Cohen (Milford, MA, USA) 11

Determination of peptides by high-performance liquid chromatography with laser-induced fluorescence detection
by H. Kanazawa, T. Nagatsuka, M. Miyazaki and Y. Matsushima (Tokyo, Japan) 23

Role of high-performance liquid chromatographic protein analysis in developing fermentation processes for recombinant human growth hormone, relaxin, antibody fragments and lymphotoxin
by F.S. Jacobson, J.T. Hanson, P.-Y. Wong, M. Mulkerrin, J. Deveney, D. Reilly and S.C. Wong (South San Francisco, CA, USA) 31

Preparative desalting of bovine serum albumin by continuous annular chromatography
by K. Reissner, A. Prior and J. Wolfgang (Götzis, Austria), H.J. Bart (Kaiserslautern, Germany) and C.H. Byers (Oak Ridge, TN, USA) 49

Ion-exchange high-performance liquid chromatographic separation of protein variants and isoforms on MCI GEL ProtEx stationary phases
by T. Adachi and H. Takayanagi (Yokohama, Japan) and A.D. Sharpe (White Plains, NY, USA) 57

Use of reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene columns for the rapid separation and purification of acid-soluble nuclear proteins
by N.Zh. Zhelev, M.J. Barratt and L.C. Mahadevan (London, UK) 65

Comparative studies on the isothermal characteristics of proteins adsorbed under batch equilibrium conditions to ion-exchange, immobilised metal ion affinity and dye affinity matrices with different ionic strength and temperature conditions
by G.M.S. Finette, Q.-M. Mao and M.T.W. Hearn (Clayton, Australia) 71

Complete structure elucidation of a globular protein by particle beam liquid chromatography–Fourier transform infrared spectrometry and electrospray liquid chromatography–mass spectrometry. Sequence and conformation of β-lactoglobulin
by V.E. Turula and R.T. Bishop (Athens, GA, USA), R.D. Ricker (Newport, DE, USA) and J.A. de Haseth (Athens, GA, USA) 91

Chiral applications

Enantiomeric resolution on Chiral-AGP with the aid of experimental design. Unusual effects of mobile phase pH and column temperature
by A. Nyström and A. Karlsson (Mölndal, Sweden) 105

Influence of chromatographic descriptors on enantioresolution of a dihydropyridine and structurally related compounds
by J. Gottfries, P. Johansson and A. Karlsson (Mölndal, Sweden) 115

Enantioseparation of fourteen O-ethyl O-phenyl N-isopropyl phosphoroamidothioates by high-performance liquid chromatography on a chiral stationary phase
by R.Y. Gao, G.S. Yang, H.Z. Yang, Z.Y. Chen and Q.S. Wang (Tianjin, China) 125

Management of nonmatrix interfering peaks in a chiral high-performance liquid chromatographic assay produced by solid-phase extraction of rat plasma
by J.R. Kagel, D.T. Rossi and C.D. Lathia (Ann Arbor, MI, USA) 129

Ruggedness of enantiomeric separation by capillary electrophoresis and high-performance liquid chromatography with methylated cyclodextrins as chiral selectors
by J. Szemán, N. Roos and K. Csabai (Budapest, Hungary) 139

Miscellaneous applications

Use of high-performance liquid chromatography with photodiode-array UV detection for the creation of a 600-compound library. Application to forensic toxicology
by Y. Gaillard and G. Pépin (Paris, France) 149

Monitoring of selected metabolites and biotransformation products from fermentation broths by high-performance liquid chromatography
by G. Morovján, G. Szakács and J. Fekete (Budapest, Hungary) 165

Separation of primary and secondary amines as their sulfonamide derivatives by reversed-phase high-performance liquid chromatography
by M.I. Saleh and F.W. Pok (Penang, Malaysia) 173

Analysis of foods for heterocyclic aromatic amine carcinogens by solid-phase extraction and high-performance liquid chromatography
by M.G. Knize, C.P. Salmon, E.C. Hopmans and J.S. Felton (Livermore, CA, USA) 179

Quantification of urinary uric acid in the presence of thymol and thimerosal by high-performance liquid chromatography
by Yu.-m. Chen, R.A. Pietrzyk and P.A. Whitson (Houston, TX, USA) 187

Quantitation of urinary nucleosides by high-performance liquid chromatography
by H.M. Liebich, C. Di Stefano, A. Wixforth and H.R. Schmid (Tübingen, Germany) 193

Use of secondary equilibria in reversed-phase high-performance liquid chromatography for the determination of dissociation constants of polyprotic leukotrienes
by J.E. Hardcastle, R. Vermillion-Salsbury, K. Zhao and I. Jano (Denton, TX, USA) 199

Glucose metabolism and bioreduction of 2-butanone by *Candida utilis* studied by means of ion-exchange chromatography
by C. Cheng and J.-H. Ma (Chungli, Taiwan) 205

Normal- and bonded-phase liquid chromatography with photodiode array detection of maytansinoids
by E. Reich (Farmville, VA, USA) and A.T. Sneden (Richmond, VA, USA) 213

High-performance liquid chromatographic determination of dehydroabietic and abietic acids in traditional Chinese medications
by B.L. Lee, D. Koh, H.Y. Ong and C.N. Ong (Singapore, Singapore) 221

Determination of herbicides by reductive amperometric detection in liquid chromatography
by N. Ruiz de Erenchun, M.A. Goicolea, Z. Gomez de Balugera, M.J. Portela and R.J. Barrio (Vitoria-Gasteiz, Spain) 227

ELECTROSEPARATION METHODS

Nonequilibrium thermodynamic separation model in capillary electrophoresis
by L. Heng (Xi'an, China), Z.-G. Wang (Beijing, China), B.-C. Lin (Dalian, China), C.-G. Xu (Xi'an, China) and R.-N. Fu (Beijing, China) 237

N-(Polyethyleneglycol monomethyl ether)-N-methylmorpholinium-based background electrolytes in capillary electrophoresis
by R.L. Williams and G. Vigh (College Station, TX, USA) 253

Chiral recognition and racemic resolution of drug enantiomers by electrophoresis based on host–guest complexation
by J. Liu, H. Coffey, D.J. Detlefsen and Y. Li (Wallingford, CT, USA) and M.S. Lee (Princeton, NJ, USA) 261

Analysis of monosaccharide composition by capillary electrophoresis
by A. Guttman (Fullerton, CA, USA) 271

Practical capillary electrophoresis method for the quantitation of the acetate counter-ion in a novel antifungal lipopeptide
by L. Zhou and A. Dovletoglou (Rahway, NJ, USA) 279

Analysis of phosphopeptides by capillary electrophoresis and matrix-assisted laser-desorption ionization–time-of-flight mass spectrometry
by Y.S. Yoo, Y.S. Han, M.J. Suh and J. Park (Seoul, Korea) 285

Identification of proteins by capillary electrophoresis–tandem mass spectrometry. Evaluation of an on-line solid-phase extraction device
by D. Figeys, A. Ducret and R. Aebersold (Seattle, WA, USA) 295

Separation of proteins and peptides by capillary electrochromatography in diol- and octadecyl-modified etched capillaries
by J.J. Pesek, M.T. Matyska and L. Mauskar (San Jose, CA, USA) 307

OTHER SEPARATION METHODS

Ultrafiltration: a technique for determining the molecular-mass distribution of group parameters of organic pollutants in natural waters
by U. Ludwig, T. Grischek, P. Neitzel and W. Nestler (Dresden, Germany) 315

Determination of molecular-mass distribution of food-grade protein hydrolyzates by size-exclusion chromatography and chemiluminescent nitrogen detection
by E.M. Fujinari (Houston, TX, USA) and J.D. Manes (Evansville, IN, USA) 323

Coupled size-exclusion chromatography–anion-exchange chromatography in the analysis of poly- and oligosaccharides
by T. Suortti (Vtt, Finland) 331

Ethanol–sodium chloride–phosphate mobile phase for size-exclusion chromatography of poly(ethylene glycol) modified proteins
by J.J. Ratto, S.R. O'Conner, A.R. Distler, G.-M. Wu, D. Hummel, M.J. Treuheit, A.C. Herman and J.M. Davis (Thousand Oaks, CA, USA) 337

Preparative separation of anthocyanins by gradient elution centrifugal partition chromatography
by J.-H. Renault, P. Thépenier, M. Zèches-Hanrot and L. Le Men-Olivier (Reims, France), A. Durand (Vitry-sur-Seine, France), A. Foucault (Reims, and Vitry-sur-Seine, France) and R. Margraff (Vitry-sur-Seine, France) 345

AUTHOR INDEX VOLS. 762 AND 763 353

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc